
Extensions of nilpotent Lie algebras of type 2

Pilar Benito and Daniel de-la-Concepción*

Universidad de la Rioja, Spain

The study and description of the structure of complex Lie algebras with nilradical a nilpotent Lie algebra of type 2 can be attacked by using $\mathfrak{sl}_2(\mathbb{C})$ -representation theory. Our results will be applied to review the classifications given in [5] and [1] of the Lie algebras with nilradical the 3-dimensional Heisenberg Lie algebra and the quasiclassical free nilpotent algebra $\mathfrak{n}_{2,3}$. The 5-dimensional Lie algebra $\mathfrak{n}_{2,3}$ is one of two free nilpotent algebras admitting an invariant metric as it is proven in [2]. According to [4], quasiclassical algebras let construct consistent Yang-Mills gauge theories.

[1] J.M. Ancochea-Bermúdez, R. Campoamor-Stursberg, L. García Vergnolle: Classification of Lie algebras with naturally graded quasi-filiform nilradicals, *J. Geometry Phys.* 61 (2011), 2168-2186.

[2] V.J. Del Barco, G.P. Ovando: Free nilpotent Lie algebras admitting ad-invariant metrics, arXiv:1104.4773v2.

[3] P. Benito, D. de-la-Concepción: On Levi extensions of nilpotent Lie algebras, *Linear Alg. Appl.* 439 (5) (2013), 1441-1457.

[4] S. Okubo: Gauge theory based upon solvable Lie algebras, *J. Phys. A: Math. Gen.* 31 (1998), 7603-7609.

[5] J.L. Rubin, P. Winternitz: Solvable Lie algebras with Heisenberg ideals, *J. Phys. A: Math. Gen.* 26 (5) (1993) 1123-1138.

[6] T. Sato: The derivations of the Lie algebras, *Tohoku. Math. Journ.*, 23 (1971) 21-36.x